

FILE COPY

## REPORT DOCUMENTATION PAGE

AD-A204 058

1b. RESTRICTIVE MARKINGS

NA

3. DISTRIBUTION / AVAILABILITY OF REPORT

Distribution Unlimited

DTIC

5. MONITORING ORGANIZATION REPORT NUMBER(S)

NA

OCT 19 1988

2. NA

2b. DECLASSIFICATION / DOWNGRADING SCHEDULE

NA

4. PERFORMING ORGANIZATION REPORT NUMBER(S)

Southern Illinois University at Carbondale

6a. NAME OF PERFORMING ORGANIZATION

Southern Illinois University  
at Carbondale

6b. OFFICE SYMBOL

(If applicable)

NA

7a. NAME OF MONITORING ORGANIZATION

Office of Naval Research

6c. ADDRESS (City, State, and ZIP Code)

Dept. of Chemistry & Biochemistry  
Carbondale, Illinois 62901-4409

7b. ADDRESS (City, State, and ZIP Code)

800 N. Quincy Street  
Arlington, VA 22217-50008a. NAME OF FUNDING / SPONSORING  
ORGANIZATION

Office of Naval Research

8b. OFFICE SYMBOL

(If applicable)

ONR

9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER

N00014-86-K-0739

8c. ADDRESS (City, State, and ZIP Code)

800 N. Quincy Street  
Arlington, VA 22217-5000

10. SOURCE OF FUNDING NUMBERS

PROGRAM  
ELEMENT NO

61153N

PROJECT  
NO

RR04106

TASK  
NO

441d-010

WORK UNIT  
ACCESSION

11. TITLE (Include Security Classification)

(U) Structure and expression of various RNAs in the Archaeobacteria

12. PERSONAL AUTHOR(S)

Gupta, Ramesh

13a. TYPE OF REPORT

Annual

13b. TIME COVERED

FROM 09/87 TO 08/88

14. DATE OF REPORT (Year, Month, Day)

88/09/05

15. PAGE COUNT

2

16. SUPPLEMENTARY NOTATION

17. COSATI CODES

FIELD	GROUP	SUB-GROUP
06	03	

18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)

Archaeobacteria, Thermophiles, tRNA, Intron

19. ABSTRACT (Continue on reverse if necessary and identify by block number)

Several transfer RNA genes of Sulfolobus solfataricus have been cloned and partially sequenced. Some of these clones are being used to determine the transcription initiation and termination sites of these genes and to study various post-transcriptional processing events.

The genes for some minor small RNAs, which migrate between 7S and 4S RNAs during polyacrylamide gel electrophoresis, have been cloned and are being characterized. In addition, facilities have been established to grow anaerobic thermophiles, e.g., Thermococcus celer, which is a strict anaerobe and grows optimally at 88°C and a pH of 5.5 to 5.8.

20. DISTRIBUTION / AVAILABILITY OF ABSTRACT

☒ UNCLASSIFIED/UNLIMITED    ☐ SAME AS RPT    ☐ DTIC USERS

21. ABSTRACT SECURITY CLASSIFICATION

(U)

22a. NAME OF RESPONSIBLE INDIVIDUAL

Dr. M. Marron

22b. TELEPHONE (Include Area Code)

202-696-4760

22c. OFFICE SYMBOL

ONR

## ANNUAL REPORT

CONTRACT NO.: N00014-86-K-0739  
R&T Code: 441d010

FULL TITLE: Structure and expression of various RNAs in the Archaeobacteria.

ABBREVIATED TITLE: RNAs in Archaeobacteria

INSTITUTION: The Board of Trustees of Southern Illinois University,  
Carbondale, IL 62901

PRINCIPAL INVESTIGATOR: Ramesh Gupta  
Chemistry and Biochemistry  
Southern Illinois University  
Carbondale, IL 62901

PERIOD OF REPORT: September 1987 - August 1988

### Summary of Project Goals:

1. Separation and sequencing of various RNAs of thermophilic archaeobacteria.
2. Sequencing of various tRNA genes of archaeobacteria (mainly thermophiles) and their surrounding regions to determine the organization of these genes and to identify the potential transcription control regions and the transcript processing sites.
3. Identification of transcription initiation and termination sites in tRNA genes of various thermophilic archaeobacteria.
4. Characterization of RNA processing in thermophilic archaeobacteria, initially using small transcripts, e.g., tRNA gene transcripts.

### Recent Accomplishments

Several HindIII fragments of Sulfolobus solfataricus DNA containing tRNA genes have been cloned in pTZ18U and 19U vectors using standard techniques. Partial restriction maps for some of these clones have been generated. Specific regions of some of these clones have been partially sequenced. Some of these tRNA genes contain introns while others do not. The sequences of the genes of interest are being confirmed by sequencing the complementary strands of DNA. Several shortened plasmids have been generated during sequencing studies, by generating deletions in the inserts of the original clones. These various clones and their deleted versions will also be used during later transcriptional and RNA processing studies. Direct sequencing of different Sulfolobus tRNAs by various RNA sequencing methods appears to be difficult due to the presence of various 2'-O-methylated residues in these RNAs. Therefore, availability of the sequences of tRNA genes will be helpful in determining the tRNA sequences.

Studies are being conducted to determine whether intron containing tRNA genes are transcribed or not. Preliminary data indicate that at least some of these genes are transcribed in the cell, as their precursor transcripts are detected in the total RNA

preparations. These studies will be continued to study the mechanisms of excision of introns and splicing of exons in these transcripts. In addition, the sites for transcription initiation and termination of some of the tRNA genes are being determined.

We have observed several specific small RNAs, which are present in extremely small quantities in S. solfataricus. These RNAs migrate between 7S and 5S and between 5S and 4S RNAs during polyacrylamide gel electrophoresis. These RNAs do not seem to contain any modified nucleosides as observed by two-dimensional thin layer chromatography of RNase T<sub>2</sub> digests of these RNAs. We are trying to determine whether these RNAs are precursors of some other stable RNAs or are products of RNA processing reactions. So far, we have not been able to directly sequence any one of these RNAs as it is difficult to isolate a specific RNA in large quantity and sufficient purity. Therefore, we are cloning the genes (DNA fragments) which hybridize to these RNAs. We have isolated some S. solfataricus clones which hybridize to these small RNAs. These clones are being characterized and sequenced.

Recently, we established the conditions to efficiently grow Thermococcus celer in serum bottles or other tightly capped bottles. The research machine shop at the University prepared a gassing manifold so that we can exchange the gases in closed systems (e.g., serum bottles) and grow anaerobic organisms. T. celer is a strict anaerobe, grows optimally at 88°C and a pH of 5.5 to 5.8. It also requires about 4% NaCl and elemental sulfur for its growth. It produces H<sub>2</sub>S and some other strong smelling substances (probably mercaptans). Its growth ceases under high pressure of H<sub>2</sub>S, which is immediately followed by lysis.

#### Plans for Next Year:

During next year, all the abovementioned works will be continued. Furthermore, since we can now grow T. celer, we shall also be using this organism to conduct studies similar to the ones mentioned before for S. solfataricus. Both of these organisms are thermophilic, yet they are phylogenetically sufficiently diverse to be classified into separate groups of archaebacteria.

SEARCHED FOR	
INDEXED	✓
DTIC TAB	
Unannounced	
Justification	
A-1	

